

Unexplained Infertility

What is it?

The diagnosis of "Unexplained Infertility" is used when no cause for the infertility can be found in either partner.

How common is this "Condition"?

Up to 1 in 10 infertile couples.

What is "Normal"?

1. Sperm Count
 - a. At least 20×10^6 per ml, 40% motile, 40% normal
 - b. If low, the test should be repeated
2. Ovulation test can be diagnosed by a combination of tests;
 - a. Temperature Chart: Ovulation can almost always be detected by analysing temperature charts
 - b. Blood Tests: By having several blood tests each week for a full cycle checks can be made on the level of progesterone produced by the ovaries
 - c. Endometrial biopsy: This is done by taking a small sample from inside the womb to check whether the lining of the womb is "in phase" with the time of the menstrual cycle. This test can be done without an anaesthetic, or as part of a full curette
3. Fallopian Tube Tests
 - a. Hysterosalpingogram: This X-Ray of the tubes can determine whether the internal shape of the uterus is normal and if the tubes are blocked.
 - b. Laparoscopy: This investigation, done under general anaesthetic, allows the doctor to visualise the tubes, ovaries and other pelvic structures
4. Hostility
 - a. Post Coital Test: At the time of ovulation, mucus is extracted from the cervix 4-6 hours after intercourse and the activity of the sperm is assessed

- b. Indirect Immuno Bead Test (IIBT): Mucus and sperm are collected and mixed with coated beads in the laboratory. If the sperm are not able to move properly and bind to the beads sperm antibodies are present

Has anything been missed?

There are some rare conditions which have been associated with unexplained infertility. However these do not always cause infertility. The tests are also often time consuming and expensive.

- Psychological Causes - (myth or reality?)
- Deficient Luteal Progesterone
- Mycoplasma infection
- Motility defects of Sperm
- Minimal Endometriosis
- Adhesions in the Uterus

How can this be treated?

1. Treat the cause
2. Try several treatments, in the hope that one may work; (i.e. "Shot gun therapy")

What is the chance of pregnancy?

If there is no abnormality found, your chance of getting pregnant without treatment after

3 years is about 1 in 3, if you have never been pregnant before.

With treatment your chances of getting pregnant may improve to 1 in 2, within 1 year.

Contact Us

Telephone: (08) 9382 2388
Facsimile: (08) 9381 3603
218 Nicholson Road, Subiaco WA 6008
concept@conceptfertility.com.au
www.conceptfertility.com.au